

Signs of Liver Problems

Liver, one of the vital body organs, is located in the thoracic region of the abdomen below the diaphragm. It regulates blood volumes and blood clotting. It plays an important role in blood purification too. It is responsible for bile production and fat and sugar metabolism. Thus, it directly and indirectly aids in development, immunization, nutrition and recreation. Usually, liver problem signs go unnoticed or are difficult to assess. They are most likely to be misinterpreted as signs of some other diseases. Therefore, liver diseases are often referred to as 'quiet killers'. If dysfunction of the liver is spotted at the initial stage, correct treatment can assist restore liver health. So, everyone needs to be aware of the early signs of liver problems.



Causes

Colon polyps are the result of abnormal cell growth. These cells do not grow and divide in an orderly way like their healthy counterparts. Instead they trigger anomalies in genes and keep dividing without the need for new cells. Small or sessile polyps are typically attached to a stalk. They are identified as adenomatous, hyperplastic, or inflammatory. Polyps in the colon can result in rectal bleeding, irritable bowel syndrome, severe pain in the abdomen, and colon cancer, if disregarded. Smoking and alcohol abuse significantly increase the risk of developing polyps of a malignant nature.

Medical Treatment for Lead Poisoning

Getting rid of the source of lead and altering diet is the primary treatment method. Secondly, chelation therapy is the most effective therapy used in treating lead poisoning and lowering the level of lead in the body.



Ibs

Colon Polyps can be Treated If Detected in Time

There are dedicated scientific procedures in place, such as colonoscopy or sigmoidoscopy that target the polyp stalk to stop further development, polypectomy or polyp elimination, and a variety of laparoscopic strategies. Another treatment is that of proctocolectomy in which the entire colon is removed. The process involves a procedure that is clinically referred to as ileal pouch-anal anastomosis. In this procedure, the surgeon constructs a pouch out of the ileum or end of the small intestine and attaches it to the rectum.

- Addition to all the given body structures, muscles and skin are also present on the lower abdomen on the left side of your body.
- Any medical condition affecting these structures can potentially cause pain in the left side of the waist.
- Sometimes, pain due to a particular condition on the right side of the abdomen, radiates to the left side.
- In such a condition, the diagnosis of the left side abdominal pain ends up being a little tough.

Disclaimer: Content in this document is an advertisement, therefore it may be biased and should not be considered an objective or independent review. Owner of this document may be compensated when you purchase a product by clicking a link in this document. The views and opinions expressed in this document are purely of the author. Any product claim or other representation about a product or service should be verified with the manufacturer, provider or party in question. Any health information in this document is for educational purposes only and is not intended to replace the advice of your health care provider.